

- depression. *Depress. Anxiety*. 2014 31, 690–698. doi: 10.1002/da.22263
35. Alagapan S, Choi KS, Heisig S, et al. Cingulate dynamics track depression recovery with deep brain stimulation. *Nature*.2023;622(7981):130-138. doi:10.1038/s41586023-06541-3
 36. Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*. 2011;106:1125–65.
 37. Smith SM, Fox PT, Miller KL, Glahn DC, Fox PM, Mackay CE, et al. Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci USA*. 2009;106:13040–5.
 38. Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-scale network dysfunction in major depressive disorder: a meta-analysis of resting-state functional connectivity. *JAMA Psychiatry*. 2015;72:603–11
 39. Cha J, Rajendra JJ, McGrath C, et al. Whole Brain Network effects of subcallosal cingulate deep brain stimulation for treatment-resistant depression. Preprint. *Res Sq*. 2023;rs.3.rs-3025802. Published 2023 Jun 14. doi:10.21203/rs.3.rs-3025802/v1
 40. Riva-Posse P, Choi KS, Holtzheimer PE, McIntyre CC, Gross RE, Chaturvedi A, Crowell AL, Garlow SJ, Rajendra JK, Mayberg HS. Defining critical white matter pathways mediating successful subcallosal cingulate deep brain stimulation for treatment-resistant depression. *Biol Psychiatry*.2014;76(12):963-9. doi:10.1016/j.biopsych.2014.03.029.